

AD-A086 377

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 20/11

AN INTEGRAL BOUND ON THE STRAIN ENERGY FOR THE TRACTION PROBLEM--ETC(11)

APR 80 J J ROSEMAN

DAA629-75-C-0024

UNCLASSIFIED

MRC-TSR-2060

ML

1 of 1

AD-A086 377

END

DATE

FILED

8-80

DTIC

② LEVEL II

MRC Technical Summary Report #2060

AN INTEGRAL BOUND ON THE STRAIN ENERGY
FOR THE TRACTION PROBLEM IN NONLINEAR
ELASTICITY WITH SMALL STRAINS

Joseph J. Roseman

ADA086377

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

April 1980

(Received February 26, 1980)

DTIC
ELECTE
JUL 11 1980
S B D

Approved for public release
Distribution unlimited

Sponsored by
U. S. Army Research
Office, Box 12211
Research Triangle Park
North Carolina 27709

80 7 7 116

UNIVERSITY OF WISCONSIN - MADISON
MATHEMATICS RESEARCH CENTER

AN INTEGRAL BOUND ON THE STRAIN ENERGY FOR THE TRACTION
PROBLEM IN NONLINEAR ELASTICITY WITH SMALL STRAINS.

Joseph J. Roseman

Technical Summary Report #2060

April 1980

ABSTRACT

For the traction boundary value problem in nonlinear elastostatics for a body which is convex in its undeformed reference state and with the assumption of sufficiently small strains (but not necessarily small displacement gradients), an upper bound is obtained for the elastic strain energy in terms of the L_2 -integral norms of the surface tractions and body forces with the constant depending only upon the ratio of the outer and inner diameters and the physical constants of the material.

This result extends previous known results in linear elasticity (infinitesimal displacement gradients) and finite elasticity (small but finite displacement gradients) into the small strain theory of nonlinear elasticity.

AMS (MOS) Subject Classifications: 73C50, 73G05

Key Words: nonlinear elastostatics, strain energy bound

Work Unit Number 2 (Physical Mathematics)

*Department of Mathematics, Tel-Aviv University, Tel-Aviv, Israel
Mathematics Research Center and Department of Mathematics,
University of Wisconsin-Madison, Madison, Wisconsin 53706

Sponsored by the United States Army under Contract Nos. DAAG29-75-C-0024,
DAAG29-80-C-0041 and DAAG29-76-G-0102.

SIGNIFICANCE AND EXPLANATION

Suppose that an elastic body is in a natural stress-free state and is then deformed by the action of applied surface tractions and internal body forces. This situation is described mathematically by a set of highly complex nonlinear partial differential equations.

In order to simplify the equations, it is often assumed that, with the possible exception of a rigid body motion, each point in the undeformed body has been displaced only an infinitesimal amount (linear elasticity) or, more generally, by a finite but small amount (finite elasticity).

However, there are many situations in which points in the body undergo relatively large displacements, all of which can not be included in a single rigid motion of the body, and yet the strains produced are relatively small. This is true for example in the case of long thin bodies.

Therefore, it is desirable to obtain results using only the assumption that the strains are small (but finite) while the displacements and displacement gradients are relatively large. In the present research, an upper bound for the energy in the body in terms of the surface tractions and body forces is obtained using only the small strain assumption whereas previous similar results had been obtained under the assumptions of linear or finite elasticity. This result is therefore applicable in many cases where the previous results were not.

Section <input checked="" type="checkbox"/>		
DDC	Buff Section <input type="checkbox"/>	
UNANNOUNCED	<input type="checkbox"/>	
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		

The responsibility for the wording and views expressed in this descriptive summary lies with MRC and not with the author of this report.

AN INTEGRAL BOUND ON THE STRAIN ENERGY FOR THE TRACTION
PROBLEM IN NONLINEAR ELASTICITY WITH SMALL STRAINS

Joseph J. Roseman*

1. INTRODUCTION.

An elastic isotropic and homogeneous body is assumed to possess a natural stress-free state in which it occupies a convex domain $D_R \subset E^3$. It is then subjected to the combined action of applied surface tractions and internal body forces and arrives at a new equilibrium state in a domain D . Using standard equations of nonlinear elastostatics and working in the context of small strain theory, we shall derive an estimate for the strain energy in the deformed body in terms of the L_2 integral norms of the surface tractions and the body forces with the constants involved depending upon the material properties and upon the geometry of the domain. The estimate is similar to one previously obtained by Bramble and Payne [1] in the context of linear elasticity and by Breuer and Roseman [2] in the context of finite elasticity.

The theory of mathematical elastostatics (cf. [3], [4]) assumes that the deformation from D_R to D can be described mathematically as a smooth one to one mapping from the domain D_R onto the domain D and that there is a relation between the geometric strain associated with the mapping and the stress in the deformed state. The fact that the deformed body is in equilibrium imposes conditions on the stress tensor which in turn leads to a set of second order nonlinear elliptic partial differential equations for the mapping vector function. These equations, together with the known conditions at the boundary, constitute an elliptic boundary value problem.

The equations of classical linear elastostatics are obtained by a linearization of this nonlinear system, a linearization which is, in effect, equivalent to treating

*Department of Mathematics, Tel-Aviv University, Tel-Aviv, Israel

Mathematics Research Center and Department of Mathematics,
University of Wisconsin-Madison, Madison, Wisconsin 53706

the derivatives of the displacement vector (the displacement being the difference between the position of a point in D and its former position in D_R) as infinitesimally small quantities. In theory, therefore, the equations of linear elasticity describe the physical model only for infinitesimally small motions, although experience has shown that they give good results in many applications.

In the theory of finite elasticity, it is assumed that the displacement gradients are sufficiently small, depending upon the geometry of the body and the material constants, but not infinitesimal. Finite elasticity should therefore provide a more accurate mathematical description of the physical model than linear elasticity and results in finite elasticity theory concerning existence, uniqueness and continuous dependence on the given data would, at the least, be valid wherever linear elasticity results are valid.

However, as Fritz John [5] points out, in practice it is possible to have small strains and relatively large displacement gradients, especially in thin rods or shells (indicating large relative "rotation" of different parts of the body). Thus, it would be especially meaningful to obtain results based on the nonlinear equations assuming only that the strains are sufficiently small (but not infinitesimal), while the displacement gradients are bounded but are permitted to be large in comparison with the strains. That is what is done in this paper.

The techniques employed here are a refinement of those used in [2] and can be extended in a straightforward manner to non-isotropic and non-homogeneous bodies. The convexity requirement on D_R is imposed in order to ensure the applicability of some work of F. John [6], [7] on the relationship between rotation and strain and can be relaxed somewhat.

Other results concerning bounds for the strain energy in elastostatics in terms of the given data include [1], [8], [9], [10] in linear elasticity, [2], [11], [12] in finite elasticity, and [13] for the displacement boundary value problem in nonlinear elasticity with small strains.* A description of some of the above mentioned results is given in [14].

*In both [13] and [14], the results in [13] are described as being within the framework of finite elasticity. However, the analysis in [13] actually requires only an a priori assumption of small strains within the elastic body.

2. PRELIMINARIES.

In the absence of surface tractions or internal body forces, an isotropic, homogeneous elastic body occupies a convex domain $D_R \subset E^3$ with inner and outer diameters d and D respectively and with a boundary, ∂D_R , which is at least smooth enough for the application of the divergence theorem. Under the action of a system of known body forces and surface tractions, D_R is mapped onto a domain D with boundary ∂D with a point (x_1, x_2, x_3) in D_R being taken to the point (y_1, y_2, y_3) in D . After deformation, the body is in static equilibrium and the mapping from D_R to D is assumed to be one to one and sufficiently smooth.

The displacement vector u_i is defined as

$$u_i = y_i - x_i \quad (2.1)$$

and, since the mapping is one to one, u_i may be regarded as a function of either (x_1, x_2, x_3) or (y_1, y_2, y_3) .

We next define*

$$p_{ik} = \frac{\partial y_i}{\partial x_k} = \delta_{ik} + \frac{\partial u_i}{\partial x_k}, \quad (2.2)$$

$$p^{ik} = \frac{\partial x_i}{\partial y_k}, \quad (2.3)$$

$$g_{ik} = p_{ji} p_{jk} = g_{ki}, \quad (2.4)$$

$$g^{ik} = p^{ij} p^{kj} = g^{ki}. \quad (2.5)$$

The tensors p_{ik} and p^{ik} are, respectively, the covariant and contravariant tensors, g_{ik} and g^{ik} are the covariant and contravariant metric tensors, and δ_{ik} is the Kronecker delta (identity tensor).

*Tensor notation is used throughout the paper and all indices may take on the values 1, 2 and 3. The summation convention is followed so that a repeated index in any term is summed over all values of the index. The magnitudes of a (real) vector v_i and matrix b_{ik} are defined here as $|v| = \sqrt{v_i v_i}$ and $|b| = \sqrt{\frac{1}{3} b_{ik} b_{ik}}$.

We have the relations

$$p^{ij}p_{jk} = p_{ij}p^{jk} = \delta_{ik} \quad (2.6)$$

and

$$g^{ij}g_{jk} = g_{ij}g^{jk} = \delta_{ik} \quad (2.7)$$

There are several definitions in the literature for the strain tensor in nonlinear elasticity. One of the more common, which we shall use here, is

$$e_{ik} = \frac{1}{2} (g_{ik} - \delta_{ik}) \quad (2.8)$$

From (2.2) and (2.4), this is equivalent to

$$e_{ik} = \frac{1}{2} \left(\frac{\partial u_i}{\partial x_k} + \frac{\partial u_k}{\partial x_i} + \frac{\partial u_j}{\partial x_i} \frac{\partial u_j}{\partial x_k} \right) \quad (2.9)$$

or, in terms of y_i coordinates,

$$e_{ik} = \frac{1}{2} \left(\frac{\partial u_i}{\partial y_k} + \frac{\partial u_k}{\partial y_i} + p_{mk} \frac{\partial u_j}{\partial y_m} \frac{\partial u_j}{\partial y_i} + p_{mi} \frac{\partial u_j}{\partial y_m} \frac{\partial u_k}{\partial y_j} + p_{ji} p_{mk} \frac{\partial u_n}{\partial y_j} \frac{\partial u_n}{\partial y_m} \right) \quad (2.10)$$

The medium is assumed to be hyperelastic, i.e. it possesses a positive definite strain energy density function $W = W(e)$ which has the properties that for every $G \subset \mathcal{D}$ whose undeformed preimage is $G_R \subset \mathcal{D}_R$, the strain energy in G , $U(G)$, is given by

$$U(G) = \iiint_{G_R} W(e) dx_1 dx_2 dx_3 \quad (2.11)$$

and

$$W(e) = \mu e_{ik} e_{ik} + \frac{\lambda}{2} e_{ii} e_{kk} + O(e^3) \quad (2.12)$$

where $O(e^3)$ is a smooth term of order of magnitude $|e|^3$ with a derivative of order $|e|^2$ for small $|e|$ and λ and μ are positive material constants.

In addition, there exist material constants m and M such that

$$m|e|^2 \leq W(e) \leq M|e|^2 \quad (2.13)$$

if $|e|$ is sufficiently small.

The stress at any point in D is represented by the Cauchy stress tensor τ_{ik} which has the property that for any differential surface element dS in D with unit normal vector v_k , the components of force on this element are $\tau_{ik} v_k dS$.

The body in the deformed state is in equilibrium under the action of the body forces per unit volume F_i and surface tractions (force/area) T_i .

The tensor τ_{ik} satisfies the equations

$$\frac{\partial \tau_{ik}}{\partial y_k} = -F_i, \quad y_i \in D, \quad (2.14a)$$

and the boundary conditions

$$\tau_{ik} n_k = T_i, \quad y_i \in \partial D, \quad (2.14b)$$

where n_i is the outward unit normal to ∂D .

The existence of the strain energy density function $W(e)$ implies the following relation between τ_{ik} and e_{ik} :

$$\tau_{ik} = \frac{1}{\det(p)} p_{km} \frac{\partial W(e)}{\partial p_{im}}, \quad (2.15a)$$

(cf. Stoker [3, pp. 10-14]) or, from (2.4) and (2.8),

$$\tau_{ik} = \frac{1}{\det(p)} p_{in} p_{km} \frac{\partial W(e)}{\partial e_{nm}}. \quad (2.15b)$$

Equation (2.15b) shows that the tensor τ_{ik} is symmetric (because e_{ik} is symmetric).

For the problem considered here, it is assumed a priori that, throughout D , $|e|$ is sufficiently small with respect to the geometry of D_R and the material constants; more precisely we assume that

$$|e| \leq \delta \left(\frac{d}{D}\right)^3 \quad (2.16)$$

with δ sufficiently small depending upon the material.

However, the only restrictions that we require on the displacement gradients are that the mapping from D_R to D be one-to-one, that the magnitudes of the matrices

p_{ik} , p^{ik} , g_{ik} and g^{ik} and their determinants be less than or equal to two in absolute value, and that \mathcal{D} satisfy some mild geometric requirements, which are stated in the theorem below.

3. THE ENERGY BOUND.

We now state the main theorem: Consider an elastic, isotropic body which possesses a stress-free reference state in which it is homogeneous and occupies a convex domain $D_R \subset E^3$ with inner and outer diameters d and D respectively. Suppose that the body is subjected to internal and external forces and reaches a state of static equilibrium in which it occupies a domain $D \subset E^3$ with internal body forces F_i (force/unit volume) throughout D and surface tractions T_i (force/unit area) on ∂D , the boundary of D .

Assume that each point (x_1, x_2, x_3) in D_R is mapped to a point (y_1, y_2, y_3) in D in a smooth, one to one manner, that the magnitudes of the covariant and contravariant Jacobian and metric matrices and the determinants of these matrices are bounded in absolute value at every point by two, and that the strain matrix satisfies (2.16).

Suppose that ∂D has continuous curvature with maximum principal curvature not greater than $32/d$ and that at every point on ∂D there exists a sphere of radius $d/64$ which is tangent to the surface at the point and whose interior lies entirely within D . In addition, there exists a point in D , which is taken to be the origin, with the properties that the interior of the sphere $\sum y_i y_i = \frac{d^2}{16}$ is contained in D and $y_i n_i / \sqrt{y_k y_k} \geq -\frac{1}{4}$ for all (y_1, y_2, y_3) on ∂D , where n_i is the outward unit normal vector.

Assume that there exists a strain energy density function for the material which satisfies the relations (2.12) and (2.13).

Then, if δ (Equation (2.16)) is sufficiently small with respect to the material constants, the total elastic energy of the body $U(D)$ satisfies the inequality

$$U(D) \leq B d D \left(\frac{D}{d}\right)^q \left[\|F\|^2 + \frac{1}{D} \|T\|^2 \right], \quad (3.1)$$

where

- i) $\|F\|$ is the L_2 integral norm of $|F|$ over D ,
- ii) $\|T\|$ is the L_2 integral norm of $|T|$ over ∂D ,
- iii) q is a sufficiently large positive universal constant, and
- iv) B is a constant which depends only upon the physical constants of the material.

The rest of Section 3 consists of the proof of this result. Throughout the following, all constants denoted by K_i are universal, while those denoted by C_i may depend also upon the material constants.

We begin by noting that, because of (2.12), (2.13) and (2.15), and because of the fact that the magnitudes and determinants of p , p^{-1} , g , and g^{-1} are uniformly bounded, that the integrals over either D_R or D of the expressions $W(e)$, $e_{ik}e_{ik}$, $e_{ik}e_{ik}$, and $\tau_{ik}e_{ik}$ are all of the same order of magnitude in the sense that any one is bounded in terms of any other with the constant depending only upon the material constants. We also see that

$$|\tau| \leq C_1 \delta \left(\frac{d}{D}\right)^3. \quad (3.2)$$

We now call attention to the fact that if the deformed body were now subjected to any rigid body motion (that is, any combination of translations and rotations), the magnitude of the strain tensor e at any point as it moves with the body would remain invariant and so would the magnitude of the stress tensor τ , the strain energy $U(\Pi)$, and the integral norms of the body forces and the surface tractions (cf. [15, pp. 127-128]). Therefore, since all the quantities of interest in (3.1) are unaffected, we shall assume that a rigid motion has already taken place and has been included in the mapping from D to D_R . The rigid motion has been chosen so that the body is in what we consider to be a convenient orientation, as will be explained below.

The papers of Fritz John [6], [7], on the relationship between rotation and strain imply that if the quantity δ in (2.16) is sufficiently small, the body may be rigidly rotated to a position in which an inequality of the following type is valid:

$$\iiint_{D_R} \frac{\partial u_i}{\partial x_k} \frac{\partial u_i}{\partial x_k} dx_1 dx_2 dx_3 \leq K_1 \frac{D^3}{d^3} \iiint_{D_R} e_{ik} e_{ik} dx_1 dx_2 dx_3. \quad (3.3)$$

Such a position is one to which we have referred as convenient in the paragraph above.

We note that the integrand on the right side of (3.3) is invariant under all translations and rotations while the integrand on the left depends upon the orientation of the body.

Because of the bounds on p and p^{-1} , we also have

$$\iiint_D \frac{\partial u_i}{\partial y_k} \frac{\partial u_i}{\partial y_k} dy_1 dy_2 dy_3 \leq K_2 \frac{D^3}{d^3} \iiint_D e_{ik} e_{ik} dy_1 dy_2 dy_3, \quad (3.4)$$

and, from the remarks in the paragraph preceding (3.2),

$$\iiint_D \frac{\partial u_i}{\partial y_k} \frac{\partial u_i}{\partial y_k} dy_1 dy_2 dy_3 \leq K_3 \frac{D^3}{d^3} \iiint_D \tau_{ik} e_{ik} dy_1 dy_2 dy_3. \quad (3.5)$$

From this point on, we shall use dv_x , ds_x , dv_y and ds_y to denote differential volume and surface elements in D_R and D .

From (2.10) and from the fact that τ_{ik} is a symmetric tensor, it follows that

$$\begin{aligned} \iiint_D \tau_{ik} e_{ik} dv_y &= \iiint_D \tau_{ik} \frac{\partial u_i}{\partial y_k} dv_y \\ &+ \iiint_D p_{mk} \tau_{ik} \frac{\partial u_j}{\partial y_m} \frac{\partial u_i}{\partial y_j} dv_y \\ &+ \frac{1}{2} \iiint_D p_{ji} p_{mk} \tau_{ik} \frac{\partial u_n}{\partial y_j} \frac{\partial u_n}{\partial y_m} dv_y. \end{aligned} \quad (3.6)$$

Using the divergence theorem and noting (2.14), we obtain

$$\begin{aligned} \iiint_D \tau_{ik} e_{ik} dv_y &= \iiint_D F_i u_i dv_y + \iint_{\partial D} T_i u_i ds_y \\ &+ \iiint_D p_{mk} \tau_{ik} \frac{\partial u_j}{\partial y_m} \frac{\partial u_i}{\partial y_j} dv_y \\ &+ \frac{1}{2} \iiint_D p_{ji} p_{mk} \tau_{ik} \frac{\partial u_n}{\partial y_j} \frac{\partial u_n}{\partial y_m} dv_y. \end{aligned} \quad (3.7)$$

Now, we have, from (3.2) and (3.5), that

$$\begin{aligned} \left| \iiint_D p_{mk} \tau_{ik} \frac{\partial u_j}{\partial y_m} \frac{\partial u_i}{\partial y_j} dv_y \right| &\leq \iiint_D |p_{mk}| |\tau_{ik}| \left| \frac{\partial u_j}{\partial y_m} \right| \left| \frac{\partial u_i}{\partial y_j} \right| dv_y \\ &\leq C_2 \left(\frac{d}{D} \right)^3 \iiint_D \frac{\partial u_i}{\partial y_k} \frac{\partial u_i}{\partial y_k} dv_y \\ &\leq C_3 \iiint_D \tau_{ik} e_{ik} dv_y. \end{aligned} \quad (3.8)$$

Similarly,

$$\left| \iiint_D p_{ji} p_{mk} \tau_{ik} \frac{\partial u_n}{\partial y_j} \frac{\partial u_n}{\partial y_m} dv_y \right| \leq C_4 \iiint_D \tau_{ik} e_{ik} dv_y. \quad (3.9)$$

Thus, for sufficiently small δ , the sum of the last two terms on the right side of (3.7) is less than $\frac{1}{2} \iiint_D \tau_{ik} e_{ik} dv_y$ and therefore

$$\frac{1}{2} \iiint_D \tau_{ik} e_{ik} dv_y \leq \iiint_D F_i u_i dv_y + \iint_{\partial D} T_i u_i dS_y. \quad (3.10)$$

Since the body is in equilibrium, the surface tractions and body forces satisfy the consistency relations (see [15], pp. 127-128)

$$\iint_{\partial D} T_i dS_y + \iiint_D F_i dv_y = 0, \quad (3.11a)$$

$$\iint_{\partial D} (y_i T_k - y_k T_i) dS_y + \iiint_D (y_i F_k - y_k F_i) dv_y = 0. \quad (3.11b)$$

The first equation above expresses the vanishing of the resultant force, while the second expresses the vanishing of the resulting moment in the deformed body.

We now define

$$v_i = u_i + \alpha_{ij} y_j + \beta_i \quad (3.12)$$

where α_{ij} is a real constant anti-symmetric tensor ($\alpha_{ik} = -\alpha_{ki}$) and β_i is a constant vector, both of which will be chosen later.

From the relations (3.11) we see that

$$\iiint_D F_i u_i dv_y + \iint_{\partial D} T_i u_i dS_y = \iiint_D F_i v_i dv_y + \iint_{\partial D} T_i v_i dS_y, \quad (3.13)$$

and then, from the Cauchy-Schwarz inequality, we obtain

$$\begin{aligned} \iiint_D F_i u_i dv_y + \iint_{\partial D} T_i u_i dS_y &\leq \left[\iiint_D F_i F_i dv_y \right]^{\frac{1}{2}} \left[\iiint_D v_i v_i dv_y \right]^{\frac{1}{2}} \\ &+ \left[\iint_{\partial D} T_i T_i dS_y \right]^{\frac{1}{2}} \left[\iint_{\partial D} v_i v_i dS_y \right]^{\frac{1}{2}}. \end{aligned} \quad (3.14)$$

Combination of (3.14) and (3.10) yields

$$\begin{aligned} \iiint_D \tau_{ik} e_{ik} dv_y &\leq 2 \left[\iiint_D F_i F_i dv_y \right]^{\frac{1}{2}} \left[\iiint_D v_i v_i dv_y \right]^{\frac{1}{2}} \\ &+ 2 \left[\iint_{\partial D} T_i T_i dS_y \right]^{\frac{1}{2}} \left[\iint_{\partial D} v_i v_i dS_y \right]^{\frac{1}{2}}. \end{aligned} \quad (3.15)$$

We now recall from the statement of the theorem that $y_i n_i / \sqrt{y_k y_k} \geq -\frac{1}{4}$ on ∂D and that the interior of the sphere $\sum y_i y_i = \frac{1}{16} d^2$ is contained in D . Next we refer to the work of Bramble and Payne [1] who proved that if the coefficients a_{ik} and β_i are chosen such that

$$\iint_{\Sigma} v_i dS_y = \iint_{\Sigma} (v_i y_j - v_j y_i) dS_y = 0, \quad (3.16)$$

then, for a sufficiently large universal constant q_1 ,

$$\iint_{\partial D} v_i v_i dS_y \leq C_5 d \left(\frac{D}{d} \right)^{q_1} \iiint_D \tilde{W}(\theta) dv_y, \quad (3.17a)$$

and

$$\iiint_D v_i v_i dv_y \leq C_6 d^2 \left(\frac{D}{d} \right)^{q_1+1} \iiint_D \tilde{W}(\theta) dv_y, \quad (3.17b)$$

where

$$\theta_{ik} = \frac{1}{2} \left(\frac{\partial v_i}{\partial y_k} + \frac{\partial v_k}{\partial y_i} \right) \quad (3.18a)$$

and

$$\tilde{W}(\theta) = \mu \theta_{ik} \theta_{ik} + \frac{1}{2} \lambda \theta_{ii} \theta_{kk}. \quad (3.18b)$$

The quantities θ_{ik} and $\tilde{W}(\theta)$ represent, respectively, the linearized strain and the corresponding linearized energy density function. We note that

$$\tilde{W}(\theta) \leq \left(\mu + \frac{3}{2} \lambda \right) \theta_{ik} \theta_{ik}. \quad (3.19)$$

From (3.12), we see that

$$\theta_{ik} = \frac{1}{2} \left(\frac{\partial u_i}{\partial y_k} + \frac{\partial u_k}{\partial y_i} \right), \quad (3.20)$$

and then, from (3.20) and (2.10),

$$\theta_{ik} = e_{ik} - \frac{1}{2} p_{mk} \frac{\partial u_j}{\partial y_m} \frac{\partial u_i}{\partial y_j} - \frac{1}{2} p_{mi} \frac{\partial u_j}{\partial y_m} \frac{\partial u_k}{\partial y_j} - \frac{1}{2} p_{ji} p_{mk} \frac{\partial u_n}{\partial y_j} \frac{\partial u_n}{\partial y_m}. \quad (3.21)$$

The elements of p_{ij} and $\frac{\partial u_i}{\partial y_j}$ are uniformly bounded and the tensor $\frac{\partial u_i}{\partial y_j}$ satisfies (3.5). Therefore,

$$\iiint_D \theta_{ik} \theta_{ik} dv_y \leq C_7 \frac{D^3}{d^3} \iiint_D \tau_{ik} e_{ik} dv_y, \quad (3.22)$$

and, from (3.19),

$$\iiint_D \tilde{w}(\theta) dv_y \leq C_8 \frac{D^3}{d^3} \iiint_D \tau_{ik} e_{ik} dv_y. \quad (3.23)$$

With $q_2 = q_1 + 3$, Equations (3.23) and (3.17) imply that

$$\iint_{\partial D} v_i v_i ds_y \leq C_9 d \left(\frac{D}{d}\right)^{q_2} \iiint_D \tau_{ik} e_{ik} dv_y, \quad (3.24a)$$

and

$$\iiint_D v_i v_i dv_y \leq C_{10} d^2 \left(\frac{D}{d}\right)^{q_2+1} \iiint_D \tau_{ik} e_{ik} dv_y. \quad (3.24b)$$

Combining (3.24) with (3.15), we obtain

$$\sqrt{\iiint_D \tau_{ik} e_{ik} dv_y} \leq 2\sqrt{C_{10}} d \left(\frac{D}{2}\right)^{\frac{q_2+1}{2}} \|F\| + 2\sqrt{C_9} d \left(\frac{D}{d}\right)^{\frac{q_2}{2}} \|T\|. \quad (3.25)$$

Finally, the remarks in the paragraph preceding (3.2) imply that there exists a constant C_{11} such that

$$U(D) \leq C_{11} \iiint_D \tau_{ik} e_{ik} dv_y, \quad (3.26)$$

and then (3.25) and (3.26) lead to the desired result, Equation (3.1).

4. SOME ADDITIONAL REMARKS.

As stated earlier in Section 1, the techniques used here are also applicable to non-isotropic and non-homogeneous bodies and the restriction here to a homogeneous isotropic body was only for the purpose of simplifying the presentation. The convexity assumption on D_R can also be relaxed somewhat; the important thing is that the domain be such that it is possible to obtain an inequality of the form (3.3) by an appropriate rotation. (See [6], [7] for a discussion of the factors involved).

ACKNOWLEDGMENT: Part of the research was done at the University of Delaware, Mathematics Department, Newark, Delaware, during the summer of 1979 and the rest during 1979-1980 while on leave of absence at the University of Wisconsin.

REFERENCES

1. J. H. Bramble and L. E. Payne, Some inequalities for vector functions with applications in elasticity. Archive for Rational Mechanics and Analysis 11, 16 (1962).
2. S. Breuer and J. J. Roseman, Integral bounds on the strain energy for the traction problem in finite elasticity. Archive for Rational Mechanics and Analysis 68, 333 (1978).
3. J. J. Stoker, Topics in Nonlinear Elasticity. New York University, Courant Institute of Mathematical Sciences Lecture Notes (1964).
4. V. V. Novozhilov, Theory of Elasticity (translated by J. K. Lusher). Pergamon Press (1961).
5. F. John, A priori estimates, geometric effects and asymptotic behavior. Bulletin of the American Mathematical Society 81, 1013 (1975).
6. F. John, Rotation and strain. Comm. Pure Appl. Math. 14, 391 (1961).
7. F. John, Bounds for deformations in terms of average strains, in Inequalities, III (Edited by O. Shisha), p. 129, Academic Press (1972).
8. J. H. Bramble and L. E. Payne, A priori bounds in the first boundary value problem in elasticity. J. of Research of the National Bureau of Standards, Section B 65B, 269 (1961).
9. A. Dou, Upper estimate of the potential elastic energy of a cylinder. Comm. Pure Appl. Math. 19, 83 (1966).
10. A. Gutierrez, A bound on the elastic potential energy of a cylinder with square cross-section. Rev. Real Acad. Ci. Exact. Fis. Natur. Madrid 70, 549 (1976).
11. M. Aron and J. J. Roseman, Integral estimates for the displacement and strain energy in nonlinear elasticity in terms of the body force. International J. of Eng. Science 15, 317 (1977).
12. S. Breuer and J. J. Roseman, A bound on the strain energy for the traction problem in finite elasticity with localized non-zero surface data. J. of Elasticity, to appear.

13. S. Breuer and J. J. Roseman, An integral bound for the strain energy in nonlinear elasticity in terms of the boundary displacements. J. of Elasticity 9, 21 (1979).
14. J. J. Roseman, Integral bounds for the strain energy in terms of surface tractions or displacements and body forces in finite elastostatics, in Transactions of the 25th Conference of Army Mathematicians, ARO Report 80-1, p. 447, U. S. Army Research Office, North Carolina (1980).
15. C. Truesdell and W. Noll, The Nonlinear Field Theories of Mechanics, in Handbuch der Physik, III/3, p. 1, Springer-Verlag (1965).

JJR:ed

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2060	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) AN INTEGRAL BOUND ON THE STRAIN ENERGY FOR THE TRACTION PROBLEM IN NONLINEAR ELASTICITY WITH SMALL STRAINS		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Joseph J. Roseman		8. CONTRACT OR GRANT NUMBER(s) DAAG29-76-G-0102 DAAG29-75-C-0024 DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 2 - Physical Mathematics
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE April 1980
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Nonlinear elastostatics, strain energy bound		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) For the traction boundary value problem in nonlinear elastostatics for a body which is convex in its undeformed reference state and with the assumption of sufficiently small strains (but not necessarily small displacement gradients), an upper bound is obtained for the elastic strain energy in terms of the L_2 -integral norms of the surface tractions and body forces with the constant depending only upon the ratio of the outer and inner diameters and the physical constants of the material. (continued)		

ABSTRACT (continued)

This result extends previous known results in linear elasticity (infinitesimal displacement gradients) and finite elasticity (small but finite displacement gradients) into the small strain theory of nonlinear elasticity.